

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 November 2001 (15.11.2001)

PCT

(10) International Publication Number
WO 01/85095 A2

(51) International Patent Classification⁷: **A61K**

(21) International Application Number: **PCT/IN01/00097**

(22) International Filing Date: **3 May 2001 (03.05.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
09/566,875 8 May 2000 (08.05.2000) US
09/640,947 17 August 2000 (17.08.2000) US
PCT/IN00/00111
09/802,793 22 November 2000 (22.11.2000) IN
9 March 2001 (09.03.2001) US

Area, Chikalthana, Aurangabad 431 210 (IN). **KHO-RAKIWALA, Habil, Fakhruddin** [IN/IN]; Wockhardt Ltd, Wockhardt Towers, Bandra-Kurla Complex, Bandra (East), Mumbai 400 051 (IN).

(74) Agent: **DE SOUZA, Noel, John**; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report
- entirely in electronic form (except for this front page) and available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(71) Applicants and

(72) Inventors: **DE SOUZA, Noel, John** [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN). **AGARWAL, Shiv, Kumar** [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN). **PATEL, Mahesh, Vitthalbhai** [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN). **BHAWSAR, Satish, Baliram** [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN). **BERI, Rupinder, Kaur** [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN). **YEOLE, Ravindra, Dattatraya** [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN). **SHETTY, Nitin** [IN/IN]; Wockhardt Research Centre, D-4, MIDC

(54) Title: **CHIRAL FLUOROQUINOLONE ARGININE SALT FORMS**

(57) Abstract: The invention relates to the new arginine salt forms of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid, a process for their preparation and pharmaceutical formulations which comprise those arginine salt forms as the active ingredient for its use in treating antimicrobial infections.

WO 01/85095 A2

CHIRAL FLUOROQUINOLONE ARGININE SALT FORMS**FIELD OF THE INVENTION**

5 The present invention relates to valuable new forms of a chiral fluoroquinolone, viz. arginine salt forms thereof, a novel process for manufacturing the novel arginine salt forms of the chiral fluoroquinolone, the use of the novel forms of the arginine salt of the chiral fluoroquinolone in the manufacture of pharmaceutical formulations and the use of the novel forms of the arginine salt of the chiral fluoroquinolone in medicine. More
10 particularly, it relates to arginine salt forms of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid, a process for preparing the same and their use in pharmaceutical formulations and medicine.

BACKGROUND OF THE INVENTION

15 The chiral fluoroquinolone known under the name S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid is described in JP Patent 63,192,753A, JP Patent 05,339,238A, and in our pending U.S. Patent Applications No. 09/566,875 and 09/640,947, WO 00/68229 and PCT Application
20 No. PCT/IN00/00111.

S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid is an optically active isomer of the racemic compound which is claimed in US Patent 4,399,134. Although S-(-)-9-fluoro-6,7-dihydro-8-(4-
25 hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid is not presently an ingredient of any commercial product, it is potentially useful as a commercial antimicrobial agent.

S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]
30 quinolizine-2-carboxylic acid has an aqueous solubility of 0.8 - 2.0 mg/ml over the pH range 8.0 - 9.5 at 28°C, thus creating problems in having to formulate the drug as a tablet or capsule, or in making formulations for gavage and parenteral injection. The need for a

salt is clearly indicated, as the lack of an appropriate salt form can hinder the development of dosage forms acceptable for systemic use in mammals.

- 5 S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid has a pKa value of 6.80 suggesting a weak acid character and thus an ability to form a salt with an appropriate base. Generally, conversion of a pharmacologically active compound into a salt form induces a change in the compound's physicochemical properties such as solubility, absorption velocity, etc. Pharmaceutically more desirable salt forms may be selected by studying whether or not a crystalline or
- 10 amorphous form, or polymorph or pseudopolymorph can be produced, and determining the properties including its physicochemical or biological properties. A pseudopolymorph is a polymorph that differs from a true polymorph by the incorporation of solvent (Solid-state Chemistry of Drugs, 2nd Ed. S.R. Byrn et al (Eds). SSCI, Inc. 1999, p-514).
- 15 Pharmaceutically acceptable salts of racemic 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid such as salts with inorganic bases and organic bases are mentioned in the text of Otsuka's U.S. Patent No.4,399,134. Besides salts with inorganic bases and organic bases, amino acid salts of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]
- 20 quinolizine-2-carboxylic acid are identified in our pending U.S. Patent Applications No. 09/566,875 and U.S. Patent Application No. 09/640,947, WO 00/68229 and PCT Application No. PCT/IN00/00111. The subject matter of these applications is incorporated herein by reference.
- 25 S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid, L-arginine salt 0.25 hydrate and S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo [Lj]quinolizine-2-carboxylic acid, L-arginine salt, 0.75 hydrate described in Examples 7 and 8 of U.S. Patent Applications 09/566,875 and 09/640,947, WO 00/68229 and PCT Application No. PCT/IN00/00111
- 30 respectively are highly hygroscopic and turn into syrups on exposure at a relative humidity of 41%. This presents major problems in bulk or manufacturing scale. Many hydrates and salts associated with water are susceptible to changes in humidity, are hygroscopic

under adverse storage conditions and during pharmaceutical processing of them to medicament forms.

Amino acid salts, inorganic base and alkali salts and organic base salts of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid were prepared and studied by the inventors and it was found that:


- (a) the arginine salt may exist in either a substantially crystalline form or a substantially amorphous form, each form having distinctive physicochemical, solubility and stability properties;
- 10 (b) the arginine salt is less prone than the sodium salt to absorb moisture at specified humidity levels;
- (c) the arginine salt, whether crystalline or amorphous, possesses a lower propensity to cause phlebitis than the sodium and potassium salts as determined in rats by intravenous administration; and
- 15 (d) the arginine salt is less toxic in rodents than the alkali salt forms.

In summary, the substantially crystalline form and substantially amorphous form of the arginine salt of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid have been found by the inventors to have very desirable properties in possessing under specified conditions, less hygroscopicity, favourable aqueous solubility, a low propensity to cause phlebitis, and favourable acute toxicity values. These forms are expected to be very useful as pharmaceutical agents as compared with the sodium salt, other inorganic base/alkali salts, organic base salts and other aminoacid salts. These advantages will be apparent from the experimental data shown hereafter.

SUMMARY OF THE INVENTION

The main objective of the invention accordingly relates to new arginine salts of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]

5 quinolizine-2-carboxylic acid of the formula I



Formula I

15 in which x denotes 0, 0.25, 0.5, 0.75, or 1.0, compositions comprising these salts, methods for preparing these salts and their use in medicine.

20 Another object of the present invention is to provide novel forms of the arginine salt of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid which can be used in full scale manufacturing of pharmaceutical formulations.

25 Still another object of the present invention is to provide a process for bulk scale production of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid arginine salts of this invention.

A further object of the present invention to provide antibacterial compositions comprising the S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt forms as an active component.

30

BRIEF DESCRIPTION OF THE DRAWINGS

The invention will now be described in further detail with reference to the accompanying drawings.

5

FIG. 1 represents the X-Ray Powder Diffraction (XRPD) spectrum of the substantially crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt of the invention.

10 FIG. 2 represents the X-Ray Powder Diffraction (XRPD) spectrum of the substantially amorphous form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt of the invention.

15 FIG. 3 represents the Differential Scanning Calorimeter (DSC) thermogram of the substantially crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt of the invention.


20 FIG. 4 represents the Differential Scanning Calorimeter (DSC) thermogram of the substantially amorphous form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt of the invention.

25 FIG. 5 shows that the particle size measured as mean mass diameter of the substantially crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt of the invention is ca. 84 μm and the particle size of the substantially amorphous form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt of the invention is ca. 32.5 μm .

30 FIG. 6 shows that the density of the substantially crystalline form of the invention is between 0.51 g/cm^3 (untapped) and 0.7 g/cm^3 (tapped), and the density of the substantially amorphous form of the invention is between 0.52 g/cm^3 (untapped) and 0.7 g/cm^3 (tapped).

DISCLOSURE OF THE INVENTION

The invention accordingly relates to new arginine salts of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid of
 5 the formula I

**Formula I**

in which x denotes 0, 0.25, 0.5, 0.75, or 1.0.

15 These salts have favourable aqueous solubility, are stable in humid conditions, have favorable bioavailability, have a lower propensity to cause phlebitis and lower toxicity when administered in mammals.

20 This invention also relates to a process for the preparation of the salts, to their use in pharmaceutical compositions and in medicine.

A preferred arginine isomer is the S-isomer, that is L-arginine.

25 A preferred arginine salt is substantially an anhydrate wherein $x = 0$. This anhydrate can exist in two polymorphic forms, one described as a substantially amorphous form (hereinafter amorphous form) and the other described as a substantially crystalline form (hereinafter crystalline form). The arginine salts of this invention are described as substantially crystalline or substantially amorphous because the degree of crystallinity of the crystalline form is not 100% and there is a degree of crystallinity in the amorphous
 30 form (see Figure 2). Each of the two forms are shown to have distinct physicochemical properties, are more stable to decomposition from uptake of moisture at specified conditions, are distinguished by increased stability, in particular during storage at

specified humidities and can be dried without caking or decomposing at elevated temperatures under reduced pressure. Both forms are particularly suitable for the preparation of stable pharmaceutical preparations, and have favourable biological properties.

5

Both forms, while less hygroscopic than the compounds of Examples 7 and 8 of U.S. Patent Applications 09/566,875 and 09/640,947, WO 00/68229 and PCT Application No. PCT/IN/00/00111, are to a certain degree hygroscopic. In stability studies at varying relative humidity conditions, both forms were found to remain stable at relative humidity values of 22% at 25°C. In stability studies at elevated temperatures, both forms remained stable with no decomposition up to 70°C, however, there was reduction in levels of water content and solvent content. On drying the forms at elevated temperatures over different periods of time and under different humidity conditions, the amounts of water found associated with the salt are altered. However, there is no change in the XRD-patterns of the forms. Without being bound to any theory, from this the inventors infer that the forms are substantially anhydrides and the values of the superficial amounts of associated water are amount due to the hygroscopic nature of the salt and not to the inventors knowledge to any incorporation of water in the lattice structure of the crystal water. The amorphous form tends to be slightly more hygroscopic than the crystalline form.

20

An anhydrate crystal form is a form that contains no solvent of crystallization (Solid-State Chemistry of Drugs, second edition, Stephen R. Byrn, Ralph R. Pfeiffer and Joseph Stowell, SSCI, Inc. West Lafayette, Indiana). Hydrates are said to exist when the drug substance incorporates water in the crystal lattice in either stoichiometric or non-stoichiometric amounts" (Stephen Byrn, et.al., Pharmaceutical Solids: A strategic Approach to Regulatory Considerations, Pharma. Res. Vol. 12 (7), 1995, 945-954).

The inventors have found that S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid arginine salt forms of this invention, have acceptable aqueous solubility of 1 mg/ml extending to 40 mg/ml, depending on the form, over the pH range 8.0 - 9.5 at ambient temperatures. The solubility of the crystalline form is 4.0 to 6.0 mg/ml at pH 9.5. The solubility of the amorphous form is 38.0

to 40.0 mg/ml at pH 9.5. The arginine salt forms are stable on heating at temperatures up to 65 – 70°C at relative humidity ranges up to 22%. They have an acceptable dissolution rate.

- 5 The combination of physical properties of the novel S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt forms of the present invention with respect to the degree of crystallinity, particle diameter, density, hygroscopicity, water content and content of other solvents are favourable and permit the manufacturing of S-(-)-9-fluoro-6,7-dihydro-8-(4-
- 10 hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt in a composition which possesses the desired properties.

The properties of the crystalline and amorphous forms including the melting points are of values that endow the forms with desirable compression and flow properties for the

15 processing of dosage forms useful for medicinal purposes.

The crystalline and amorphous forms of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt also have desirable physicochemical properties as hereinbefore mentioned, and are characterised by

20 parameters such as XRD, DSC, particle size and powder density.

The substantially crystalline form is described below:

- (a) The degree of crystallinity as determined by X-ray powder diffraction is shown in Fig. 1, X-ray powder diffraction (2 θ): 10.16, 11.78, 12.52, 16.00, 18.94, 19.66, 20.36, 21.28,
- 25 21.92, 22.52, 24.74, 25.28, 30.74.
- (b) A thermogram as determined by Differential scanning calorimetry is shown in Fig. 3 (DSC exotherm at 242.18°C (onset at 207.85°C)).

It is desirable that the substantially crystalline form also has the following properties:

30

- (c) Particle size measured as mean mass diameter (MMD) less than 99 μ m, preferably less than 85 μ m as determined by laser diffraction technique, and as shown in Fig. 5;

- (d) Density between 0.51 g/cm³ (untapped) and 0.7 g/cm³ (tapped) As shown in Fig. 6;
- (e) Hygroscopicity not exceeding 0.1 % increase of weight upon storage for 14 days up to 22% relative atmospheric humidity as determined gravimetrically;
- (f) A content of water between 0.1 and 0.2% by weight as determined by titration according to Karl Fischer; and
- (g) When methyl ethylketone, methylisobutylketone or acetone is used in the preparation of this form, the content of the solvent is less than 0.5 %, preferably less than 0.4 % by weight as determined by gas chromatography. When acetonitrile is used as solvent, the content of acetonitrile is less than 0.04 %, preferably less than 0.03 % by weight as determined by gas chromatography.

The substantially amorphous form is described below:

- (a) The degree of crystallinity as determined by X-ray powder diffraction is shown in Fig. 2, X-ray powder diffraction (2 θ): 18.28, 18.8, 19.8, 20.12, 20.62, 21.10, 21.44, 21.88, 22.6, 23.02.
- (b) A thermogram as determined by Differential scanning calorimetry is shown in Fig. 4 (DSC endotherm at 194.60°C (onset at 185.28°C) and an exotherm at 245.81°C (onset at 226.23°C)).
- It is desirable that the substantially amorphous form also has the following properties:
- (c) Particle size measured as mean mass diameter (MMD) less than 43 μ m, preferably less than 34 μ m as determined by laser diffraction technique, and as shown in Fig. 5;
- (d) Density between 0.52 g/cm³ (untapped) and 0.7 g/cm³ (tapped) as shown in Fig. 6;
- (e) Hygroscopicity not exceeding 0.1 % increase of weight upon storage for 14 days up to 22% relative atmospheric humidity as determined gravimetrically;
- (f) A content of water between 0.4 and 0.7% by weight as determined by titration according to Karl Fischer; and
- (g) When an alkanol is used as the solvent in the preparation of the substantially amorphous form, the content of solvent is less than 0.3% by weight, preferably less than 0.2% by weight as determined by gas chromatography.

The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo

[i,j]quinolizine-2-carboxylic acid arginine salt exhibits the same antibacterial activity as the free active ingredient, S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid.

- 5 The substantially crystalline and substantially amorphous forms of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]-quinolizine-2-carboxylic acid arginine salt anhydrate have antimicrobial activity:**

They have potent activity against resistant *Staphylococcus aureus*.

10

They have ability to resist drug efflux in Gram-positive organisms.

They have the unusual ability to retain potency at an acidic pH of 5.5.

- 15 They are effective in treating respiratory pathogens.**

The are effective in treating resistant mutants.

They have superior cidal action against slow-growing *Staphylococci*.

20

They score over the standard drugs in terms of eradication efficacy for *Staphylococci* from vital organs and thigh muscle.

- 25 The present invention also relates to a process for preparing the novel polymorphic forms of the arginine salt. A process for the manufacture of the substantially crystalline and substantially amorphous forms of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt comprises the following consecutive steps:**

- 30 a) treating S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo[i,j]quinolizine-2-carboxylic acid with arginine in a solvent mixture composed of water and an organic solvent, to form a solution;**

- b) concentrating the solution under reduced pressure to provide a residue;
 - c) treating the residue with an alkane;
 - d) isolating the substantially crystalline or substantially amorphous form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt; and
 - e) purifying and drying the substantially crystalline or substantially amorphous form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo [i,j] quinolizine-2-carboxylic acid arginine salt using conventional methods.
- 10 Preferably the ratio of arginine to S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H, benzo [i,j] quinolizine-2-carboxylic acid is 1 mole to 1 mole.

Preferably the solvent mixture used in step (a) is composed of water and an organic solvent. When the organic solvent is a ketone or nitrile, the content of water in the solvent mixture is 20% - 50%. When the organic solvent is methanol, the content of water in the solvent mixture is 2% - 10%; when the organic solvent is ethanol, the water content in the solvent mixture is about 15%; when the organic solvent is isopropanol, the water content in the solvent mixture is about 50%.

- 20 Preferably the solution of step (a) is stirred for 30-60 minutes.

Preferably in step (b) the solution is concentrated at 170-180 m bar and then at 40-60 m bar.

- 25 In a preferred embodiment, the aqueous solution of arginine is added to a solution of an equimolar amount of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid in an organic solvent, stirred for 30 - 60 minutes, and concentrated under reduced pressure, first at 170 -180 mbar and then at 40 - 60 mbar to provide a residue. Organic solvents may be selected from (C₁-C₆) ketones, (C₁-C₆) nitriles, or (C₁-C₆) alkanols and the like. When the organic solvent used is a ketone or nitrile such as methylethylketone, methylisobutylketone, methylbutylketone, acetone or acetonitrile, the residue obtained is substantially crystalline. When the organic solvent
- 30

used is one such as an alkanol, for example methanol, ethanol or isopropanol, the residue obtained is substantially amorphous. The residue is treated with an alkane such as a (C₁-C₆) alkane, preferably hexane or pentane. The mixture is stirred for 3 – 5 hours, and the desired salt isolated by filtration or centrifugation, washed if necessary with additional amounts of alkane, preferably hexane or pentane, and dried preferably under reduced pressure and heating up to 65 – 70° C.

The process for manufacturing the new forms of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt differs from the earlier known processes in that it can be controlled to give two different polymorphic forms of the substantially anhydrate salt form through use of the respective solvents that are shown to result in the two forms. This process can be carried out in conventional chemical process equipment and produces a product, which is substantially an anhydrate. It was not possible to obtain these products with known processes.

A comparison between the different samples of the novel forms of the arginine salt prepared by the process of the present invention and those obtained from the experiments disclosed in the prior art shows that the salt of the present invention is substantially an anhydrate, is stable up to 70°C and at relative humidity up to 22% at 25°C, and has favourable aqueous solubility, all attributes essential to the preparation of pharmaceutical compositions. The novel forms of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt are also stable under typical storage conditions, have good bioavailability in mammals, have lower phlebitis-forming potential on administration to mammals, have low or reduced toxicity, have acceptable disintegration and dissolution rates, and hence are very useful for pharmaceutical manufacturing and for use in medicine. The forms are specially suitable for long-term intravenous therapy in critically ill patients or patients in intensive care units. Injectable preparations of the arginine salt can be readily prepared in view of its availability in a bulk form that remains stable under specified conditions, its favourable aqueous solubility, its ideal suitability in not causing venous inflammation on repeated intravenous administration, and its safety from adverse toxicity.

An **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt** form of this invention can be present in pharmaceutical formulations as the only active compound or can be combined with other active ingredients such as other antibacterial agents.

5

The invention therefore also relates to liquid and solid pharmaceutical formulations which comprise the arginine salt of the invention, such as for example, injectable solutions, suspensions, emulsions, tablets, coated tablets, coated tablet cores, capsules, solutions, troches, dispersions, patches, powders, lotions, gels, sprays, pellets, granules, suppositories, hard or soft gelatin capsules, ointments, creams and the like.

10

The pharmaceutical formulations are prepared in a manner known per se, for example by mixing, stirring, suspending, dispersing, emulsifying, dissolving and the like, the active compounds with or in the pharmaceutical auxiliaries such as a carrier, diluent, solvent or excipient and processing the components to pharmaceutically suitable forms for parenteral, oral, topical, intranasal, buccal or rectal administration and the like.

15

Pharmaceutical formulations can be formulated together with auxiliaries and additives usually employed in pharmacy, such as tablet binders, fillers, preservatives, tablet disintegrating agents, flow regulating agents, plasticizers, wetting agents, dispersing agents, emulsifiers, solvents, pH altering additives, flavourings and the like.

20

The total daily dose range is generally from about 200mg to about 1500mg of the arginine salt form. However, the dose may be higher or lower depending on the needs and conditions of the patient.

25

The following detailed examples serve to more fully illustrate the invention without limiting its scope.

30 EXAMPLE 1

S-(-)-9-Fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt **Synthesis of Substantially Crystalline product**

A solution of L-(+)-arginine (48.372 g, 0.278 mole) in distilled water (600 ml) was added dropwise over a period of 30 min to the stirred solution/suspension of finely powdered S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid (100 g, 0.278 mole) in acetone (1250 ml). The obtained clear solution was stirred for 30 min and concentrated on a water bath in vacuum (175 mbar) at 80°C. When product started solidifying, the concentration was carried out in vacuum (50 mbar) at 80°C up to dryness. Hexane (1 liter) was added, the reaction mixture was stirred for 4 hr, the solid thus separated was filtered and dried in vacuum (0.7 mbar) for 12 hrs at 70 °C. Yield 145 g (96.9%), m.p. 238-242 °C, and solubility 6 mg/ml (pH 9.5 buffer solution).

The substantially crystalline S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt prepared according to Example 1 possesses the following properties:

- a) Crystalline form, with a degree of crystallinity as determined by X-ray powder diffraction and as shown in Fig. 1.
- b) A thermogram as determined by Differential scanning calorimetry and as shown in Fig. 3.
- c) Particle size measured as mean mass diameter (MMD) of 83.92 µm, as determined by laser diffraction technique.
- d) Density of 0.51 g/cm³ (untapped) and 0.7 g/cm³ (tapped).
- e) Hygroscopicity of 0% increase of weight upon storage for 14 days up to 22% relative atmospheric humidity as determined gravimetrically.
- f) A content of moisture water of 0.1 % by weight as determined by titration according to Karl Fischer.
- g) A content of acetone of 0.014 % by weight as determined by gas chromatography.

EXAMPLE 2

Similarly, when acetonitrile was used in place of acetone in the process described in Example 1, and the amount of water in the resulting acetonitrile-water solution was about 40%, a substantially crystalline product was obtained. Yield 87%.

EXAMPLE 3**Synthesis of Substantially Amorphous product**

When methanol was used in place of acetone in the process described in Example 1, and
5 the amount of water in the resulting methanol-water solution was between 2% to 10%, a
substantially amorphous product was obtained. Yield 97.3 %, m.p. 238 - 244 °C, $[\alpha]_D^{25} -$
166.2 ° (c = 1, methanol), and solubility 40 mg/ml in pH 9.5 buffer solution.

The novel form of the substantially amorphous S-(-)-9-fluoro-6,7-dihydro-8-(4-
10 hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid
arginine salt according to Example 3 possesses the following properties:

- a) Amorphous form, with a degree of crystallinity as determined by X-ray powder
diffraction and as shown in Fig. 2.
- b) A thermogram as determined by Differential scanning calorimetry and shown in Fig. 4.
- 15 c) Particle size measured as mean mass diameter (MMD) of 32.44 μm , as determined by
laser diffraction technique.
- d) Density of 0.52 g/cm³ (untapped) and 0.7 g/cm³ (tapped).
- e) Hygroscopicity of 0% increase of weight upon storage for 14 days up to 22% relative
atmospheric humidity as determined gravimetrically.
- 20 f) A content of moisture water of 0.41 % by weight as determined by titration according
to Karl Fischer.
- g) A content of methanol of 0.16 % by weight as determined by gas chromatography.

EXAMPLE 4

25

Substantially amorphous product was also obtained, when ethanol was used in place of
acetone in the process described in Example 1 and the amount of water in the resulting
ethanol-water solution was 13 – 15 %. Yield 97.44 %.

EXAMPLE 5

30

Similarly, when isopropanol was used in place of acetone as described in the process of

Example 1, and the amount of water in the resulting isopropanol-water solution was about 50%, a substantially amorphous product was obtained. Yield 80 %.

EXAMPLE 6

5

TEST EXAMPLE

X-ray Diffraction Analysis

300 mg each of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [L,j]quinolizine-2-carboxylic acid L-(+)-arginine salt prepared as in Examples 1 and 5 were thinly spread on a sample holder. X-ray diffraction analyses (40kv x 40 mA Rigaku D/max 2200) were performed under the conditions listed below:

Scan speed 5°/ min
Sampling time 7 min
Scan mode: continuous
2θ/θ reflection
Cu target (Ni filter)

Results of the X-ray diffraction analysis on the substantially crystalline and substantially amorphous forms are depicted in Fig. 1 and Fig. 2 respectively. From these spectra it can be verified that their crystal forms differ from each other.

EXAMPLE 7

TEST EXAMPLE

25 Thermal Analysis of the S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [L,j]quinolizine-2-carboxylic acid L-(+)-arginine salts prepared as in Examples 1 and 5.

For the Differential Scanning Calorimetry, METTLER TOLEDO STAR system was used. 5.4 mg of the sample was weighed into the aluminum pan, which was then press sealed with an aluminium lid. After three tiny needle holes were made on the lid the sample was tested by heating from (30°C) to (300°C) at a rate of 10°C/min. As can be seen from Fig. 4

(amorphous form) there is an endothermic peak which begins at around 185°C to 200°C and an exothermic peak due to thermal decomposition at around 226°C to 251°C. In contrast the substantially crystalline form (Fig. 3) shows only an exothermic peak at around 208°C to 256°C without any endothermic peak.

5

EXAMPLE 8**TEST EXAMPLE****BULK DENSITY DETERMINATION**

10 For untapped and tapped density determination Bulk Density Apparatus (R.V. Electronics, Mumbai) was used. 25 g of the sample was slowly poured into a dry clean stoppered measuring cylinder and filled volume was measured to obtain untapped density.

Untapped density = 25 g/untapped volume.

15

Then the measuring cylinder was fixed to Bulk Density Apparatus and sample was tapped (150 times) and volume was measured to furnish the tapped volume.

Tapped density = 25 g/Tapped volume.

20 **EXAMPLE 9**

TEST EXAMPLE**MEAN MASS DETERMINATION**

For mean mass diameter Master Sizer of Malvern Instrument LTD U.K. was used under
25 the conditions listed below:

Lens/Focus: 300 mm

Solvent: Hexane

Analysis model: Polydisperse

Obstruction value: 20-23%


30

Results are depicted in Fig. 5.

CLAIMS

1. **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt of the formula I,**

5



10

wherein x is 0, 0.25, 0.50, 0.75, or 1.0, in either a substantially crystalline form or a substantially amorphous form.

2. A **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt according to claim 1, wherein $x = 0$.**
3. A **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt having the following X-ray powder diffraction data: (2 θ): 10.16, 11.78, 12.52, 16.00, 18.94, 19.66, 20.36, 21.28, 21.92, 22.52, 24.74, 25.28, 30.74.**
4. A **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt having the following X-ray powder diffraction data: (2 θ): 10.16, 11.78, 12.52, 16.00, 18.94, 19.66, 20.36, 21.28, 21.92, 22.52, 24.74, 25.28, 30.74; a DSC exotherm at 242.18°C (onset at 207.85°C) and a water content of between 0.1 and 0.2% by weight as determined by titration according to Karl Fischer.**
5. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt having the following X-ray powder diffraction data : (2 θ): 18.28, 18.8, 19.8, 20.12, 20.62, 21.10, 21.44, 21.88, 22.6, 23.02.**

30

6. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt having the following X-ray powder diffraction data : (2 θ): 18.28, 18.8, 19.8, 20.12, 20.62, 21.10, 21.44, 21.88, 22.6, 23.02; having a DSC endotherm at 194.60°C (onset at 185.28°C) and an exotherm at 245.81°C (onset at 226.23°C), and a water content of between 0.4 and 0.7% by weight as determined by titration according to Karl Fischer.
7. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt according to claim 3, having an DSC exotherm at 242.18°C (onset at 207.85°).
8. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt according to claim 5, having a DSC endotherm at 194.60°C (onset at 185.28°C) and an exotherm at 245.81°C (onset at 226.23°C).
9. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt according to claim 3, wherein the solubility in a solution of pH 9.5 is 4.0 to 6.0 mg/ml.
10. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 4, wherein the solubility in a solution of pH 9.5 is 4.0 to 6.0 mg/ml.
11. S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt according to claim 5, wherein the solubility in a solution of pH 9.5 is 38.0 to 40.0 mg/ml.
12. S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt according to claim 6, wherein the solubility in a solution of pH 9.5 is 38.0 to 40.0 mg/ml.

13. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 3, wherein the mean mass diameter as determined by laser diffraction technique is less than 99 μm .
- 5 14. The arginine salt according to claim 13, wherein the diameter is less than 85 μm .
15. The arginine salt according to claim 5, wherein the mean mass diameter is for the substantially amorphous form is less than 43 μm .
- 10 16. The salt of claim 15, wherein the mean mass diameter is less than 34 μm .
17. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 3, wherein the density is between 0.51 g/cm^3 (untapped) and 0.7 g/cm^3 (tapped).
- 15 18. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt according to claim 4, wherein the density is between 0.51 g/cm^3 (untapped) and 0.7 g/cm^3 (tapped).
- 20 19. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxy piperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i ,j]quinolizine-2-carboxylic acid arginine salt according to claim 5, wherein the density is between 0.52 g/cm^3 (untapped) and 0.7 g/cm^3 (tapped).
20. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxy piperidin-1-yl)-5-methyl-1-oxo-1H,5H-
25 benzo [i ,j]quinolizine-2-carboxylic acid arginine salt according to claim 6, wherein the density is between 0.52 g/cm^3 (untapped) and 0.7 g/cm^3 (tapped).
21. The S-(-)-9- fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 1, wherein the
30 water content is between 0.1 % and 0.7 % by weight as determined by titration according to Karl Fischer.

22. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt** according to claim 3, wherein the water content is between 0.1 and 0.2% by weight as determined by titration according to Karl Fischer.
- 5
23. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt** according to claim 5, wherein the water content is between 0.4 and 0.7% by weight as determined by titration according to Karl Fischer.
- 10
24. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt** according to claim 3, having a solvent content less than 0.5 % by weight as determined by gas chromatography.
- 15
25. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt** according to claim 4, having a solvent content less than 0.5 % by weight as determined by gas chromatography.
- 20
26. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt** according to claim 5, having a solvent content less than 0.3% by weight as determined by gas chromatography.
- 25
27. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt** according to claim 6, having a solvent content less than 0.3% by weight as determined by gas chromatography.
- 30
28. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt** according to claim 1, wherein the hygroscopicity is less than 0.1 % increase of weight upon storage for 14 days at up to 22% relative atmospheric humidity as determined by gravimetry.
29. A process for the manufacture of **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-**

yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 1 comprising the steps of:

- 5 (a) mixing S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo[i,j]quinolizine-2-carboxylic acid with arginine in a solvent mixture comprising water and an organic solvent, to obtain a solution,
- (b) concentrating the solution under reduced pressure, to obtain a residue,
- (c) treating the residue with an alkane, and
- 10 (d) isolating S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt.
30. A process according to claim 29, wherein the amounts used of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid and arginine are equal molar amounts.
- 15 31. A process according to claim 29, wherein the organic solvent is selected from (C₁-C₆)ketones, (C₁-C₆)nitriles, and (C₁-C₆)alkanols.
32. The process according to claim 31, wherein the solvent is acetone, methylethylketone, 20 methylisobutylketone, methylbutylketone or acetonitrile.
33. The process according to claim 31, wherein the solvent is methanol, ethanol or isopropanol.
- 25 34. A process according to claim 29, wherein the alkane is hexane or pentane.
35. A process for preparing S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 3, which comprises the steps of:
- 30 (a) mixing S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo[i,j]quinolizine-2-carboxylic acid with arginine in a solvent mixture comprising water and an organic solvent selected from (C₁-C₆)ketones or (C₁-C₆)nitriles, to obtain a

solution,

- (b) concentrating the solution under reduced pressure, to obtain a residue,
- (c) treating the residue with an alkane, and
- (d) isolating the substantially crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt.

36. A process for preparing S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 5, which comprises the steps of:

- (a) mixing S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo[i,j]quinolizine-2-carboxylic acid with arginine in a solvent mixture comprising water and an organic solvent selected from (C₁-C₆)alkanols, to obtain a solution,
- (b) concentrating the solution under reduced pressure, to obtain a residue, treating the residue with an alkane, and
- (c) isolating the substantially crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt.

20

37. The process according to claim 35, wherein the percentage of water in the solvent mixture is 20% - 50%.

38. The process according to claim 36, wherein the percentage of water in the solvent mixture is 2% to 10% when the solvent is methanol, about 15% when the solvent is ethanol and about 50% when the solvent is isopropanol.

39. The process according to claim 29, wherein the isolation of the S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt is performed by centrifugal separation of the salt

40. The process according to claim 39, wherein the isolation of the S-(-)-9-fluoro-6,7-

dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid arginine salt is performed by filtration of the salt.

- 5 41. The process according to claim 29, further comprising the steps of purifying and drying the S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt.
- 10 42. The process according to claim 41, wherein the purification of the S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid arginine salt comprises by washing the crystals with a solution of alkane solvents.
- 15 43. The process according to claim 41, wherein the drying of the S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid arginine salt comprises evaporating the remaining solvent by heating.
- 20 44. A composition comprising S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 1 and a carrier, diluent, solvent or excipient.
- 25 45. A composition comprising S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 2 and a carrier, diluent, solvent or excipient.
- 30 46. A composition comprising S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 3 and a carrier, diluent, solvent or excipient.
47. A composition comprising S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 4 and a carrier, diluent, solvent or excipient.

48. A composition comprising S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 5 and a carrier, diluent, solvent or excipient.
- 5 49. A composition comprising S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 6 and a carrier, diluent, solvent or excipient.
- 10 50. A method for treating a disease caused by a microbial infection in a mammal comprising administering an effective amount of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 1 to a host in need thereof.
- 15 51. A method for treating a disease caused by a microbial infection in a mammal comprising administering an effective amount of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 2, to a host in need thereof.
- 20 52. A method for treating a disease caused by a microbial infection in a mammal comprising administering an effective amount of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 3, to a host in need thereof.
- 25 53. A method for treating a disease caused by a microbial infection in a mammal comprising administering an effective amount of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 4, to a host in need thereof.
- 30 54. A method for treating a disease caused by a microbial infection in a mammal comprising administering an effective amount of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 5, to a host in need thereof.

55. A method for treating a disease caused by a microbial infection in a mammal comprising administering an effective amount of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt according to claim 6, to a host in need thereof.


NDD/PCT/005

FIG.1

Sample : SAMPLE 1
Comment :File : SAMP 130
Memo :

Date : 19-Dec-00 19:55

Operator : RINT



NDD/PCT/005


FIG. 2

Sample : SAMPLE 2
Comment :

File : SAMP 129
Memo :


Date : 19-Dec-00 19:36

Operator : RINT




NDD/PCT/005

FIG. 3.



NDD/PCT/005


FIG. 4



METTLER TOLEDO STAR® System


NDD/PCT/005

FIG. 5



NDD/PCT/005

FIG. 6



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 November 2001 (15.11.2001)

PCT

(10) International Publication Number
WO 01/085095 A3

(51) International Patent Classification⁷: **C07D 455/04**,
C07C 279/14, A61K 31/4375, A61P 31/00

(21) International Application Number: **PCT/IN01/00097**

(22) International Filing Date: **3 May 2001 (03.05.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
09/566,875 **8 May 2000 (08.05.2000)** **US**
09/640,947 **17 August 2000 (17.08.2000)** **US**
PCT/IN00/00111
09/802,793 **22 November 2000 (22.11.2000)** **IN**
9 March 2001 (09.03.2001) **US**

(71) Applicant (for all designated States except US): **WOCKHARDT LIMITED** [IN/IN]; Wockhardt Towers, Bandra Kurla Complex, Bandra (East), Mumbai 400 051 (IN).

(71) Applicants and

(72) Inventors: **DE SOUZA, Noel, John** [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN). **AGARWAL, Shiv, Kumar** [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN). **PATEL, Mahesh, Vithalbhai** [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN). **BHAWARSAR, Satish, Baliram** [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN). **BERI, Rupinder, Kaur** [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN). **YEOLE, Ravindra, Dattatraya** [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN). **SHETTY,**

Nitin [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN). **KHO-RAKIWALA, Habil, Fakhruddin** [IN/IN]; Wockhardt Ltd, Wockhardt Towers, Bandra-Kurla Complex, Bandra (East), Mumbai 400 051 (IN).

(74) Agent: **DE SOUZA, Noel, John**; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210 (IN).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
3 October 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/085095 A3

(54) Title: **CHIRAL FLUOROQUINOLIZINONE ARGININE SALT FORMS**

(57) Abstract: The invention relates to the new arginine salt forms of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid, a process for their preparation and pharmaceutical formulations which comprise those arginine salt forms as the active ingredient for its use in treating antimicrobial infections.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IN 01/00097

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D455/04 C07C279/14 A61K31/4375 A61P31/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 399 134 A (NAKAGAWA KAZUYUKI ET AL) 16 August 1983 (1983-08-16) cited in the application claims 1,23; example 1 ----	1,50
P,X	WO 00 68229 A (PATEL M V (IN)) 16 November 2000 (2000-11-16) cited in the application claims 1,6,16; examples 8,9 -----	1,50



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

15 July 2002

Date of mailing of the international search report

26/07/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Alfaro Faus, I

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IN 01/00097

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 50-55 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

nation on patent family members

International Application No

PCT/IN 01/00097

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4399134	A	16-08-1983	
		JP 1601029 C	31-01-1991
		JP 2022074 B	17-05-1990
		JP 57176987 A	30-10-1982
		JP 1007996 B	10-02-1989
		JP 1525740 C	30-10-1989
		JP 57081486 A	21-05-1982
		AT 393383 B	10-10-1991
		AT 288789 A	15-03-1991
		AT 395150 B	25-09-1992
		AT 474881 A	15-02-1992
		AU 546358 B2	29-08-1985
		AU 7733581 A	05-08-1982
		CA 1179341 A1	11-12-1984
		CH 648845 A5	15-04-1985
		DE 3144455 A1	24-06-1982
		DE 3153221 C2	04-07-1991
		DK 495281 A ,B,	11-05-1982
		ES 507493 D0	01-02-1983
		ES 8301981 A1	01-04-1983
		ES 516523 D0	01-12-1983
		ES 8401035 A1	16-02-1984
		FI 813526 A ,B,	11-05-1982
		FR 2493849 A1	14-05-1982
		GB 2086905 A ,B	19-05-1982
		IT 1171646 B	10-06-1987
		KR 8801625 B1	02-09-1988
		MX 155102 A	27-01-1988
		NL 8105075 A ,B,	01-06-1982
		NO 813726 A ,B,	11-05-1982
		PH 16685 A	13-12-1983
		PT 73957 A ,B	01-12-1981
		SE 448542 B	02-03-1987
		SE 8106642 A	14-06-1982
		SU 1366055 A3	07-01-1988
		SU 1277896 A3	15-12-1986
		US 4552879 A	12-11-1985
		BE 891046 A1	01-03-1982
		ZA 8107733 A	27-10-1982
WO 0068229	A	16-11-2000	
		AU 5425600 A	21-11-2000
		EP 1175217 A2	30-01-2002
		WO 0068229 A2	16-11-2000

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 November 2001 (15.11.2001)

PCT

(10) International Publication Number
WO 01/085095 A3

(51) International Patent Classification⁷: **C07D 455/04**,
C07C 279/14, A61K 31/4375, A61P 31/00

Chikalthana, Aurangabad 431 210 (IN). SHETTY, Nitin
[IN/IN]; Wockhardt Research Centre, D-4, MIDC Area,
Chikalthana, Aurangabad 431 210 (IN).

(21) International Application Number: PCT/IN01/00097

(22) International Filing Date: 3 May 2001 (03.05.2001)

(74) Agent: **DE SOUZA, Noel, John**; Wockhardt Research
Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431
210 (IN).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

09/566,875	8 May 2000 (08.05.2000)	US
09/640,947	17 August 2000 (17.08.2000)	US
PCT/IN00/00111		
	22 November 2000 (22.11.2000)	IN
09/802,793	9 March 2001 (09.03.2001)	US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GI, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **WOCK-
HARDT LIMITED** [IN/IN]; Wockhardt Towers, Bandra
Kurla Complex, Bandra (East), Mumbai 400 051 (IN).

(71) Applicants and

(72) Inventors: **DE SOUZA, Noel, John** [IN/IN]; Wock-
hardt Research Centre, D-4, MIDC Area, Chikalthana,
Aurangabad 431 210 (IN). **AGARWAL, Shiv, Kumar**
[IN/IN]; Wockhardt Research Centre, D-4, MIDC Area,
Chikalthana, Aurangabad 431 210 (IN). **PATEL, Mahesh,
Vithalbhai** [IN/IN]; Wockhardt Research Centre, D-4,
MIDC Area, Chikalthana, Aurangabad 431 210 (IN).
BHAWSAR, Satish, Baliram [IN/IN]; Wockhardt Re-
search Centre, D-4, MIDC Area, Chikalthana, Aurangabad
431 210 (IN). **BERI, Rupinder, Kaur** [IN/IN]; Wockhardt
Research Centre, D-4, MIDC Area, Chikalthana, Au-
rangabad 431 210 (IN). **YEOLE, Ravindra, Dattatraya**
[IN/IN]; Wockhardt Research Centre, D-4, MIDC Area,

Published:

- with international search report
- with amended claims

(88) Date of publication of the international search report:
3 October 2002

Date of publication of the amended claims: 27 December 2002

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*


(54) Title: CHIRAL FLUOROQUINOLIZINONE ARGININE SALT FORMS

(57) Abstract: The invention relates to the new arginine salt forms of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[*l,j*]quinolizine-2-carboxylic acid, a process for their preparation and pharmaceutical formulations which comprise those arginine salt forms as the active ingredient for its use in treating antimicrobial infections.

AMENDED CLAIMS

[received by the International Bureau on 23 August 2002 (23.08.02);
original claims 4, 6-8 amended; remaining claims unchanged (2 pages)]

1. **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt of the formula I,**



wherein x is 0, 0.25, 0.50, 0.75, or 1.0, in either a substantially crystalline form or a substantially amorphous form.

2. A **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt according to claim 1, wherein $x = 0$.**
3. A **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt having the following X-ray powder diffraction data: (2θ): 10.16, 11.78, 12.52, 16.00, 18.94, 19.66, 20.36, 21.28, 21.92, 22.52, 24.74, 25.28, 30.74.**
4. A **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt having the following X-ray powder diffraction data: (2θ): 10.16, 11.78, 12.52, 16.00, 18.94, 19.66, 20.36, 21.28, 21.92, 22.52, 24.74, 25.28, 30.74; a DSC endotherm at 242.18°C (onset at 207.85°C) and a water content of between 0.1 and 0.2% by weight as determined by titration according to Karl Fischer.**
5. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt having the following X-ray powder diffraction data : (2θ): 18.28, 18.8, 19.8, 20.12, 20.62, 21.10, 21.44, 21.88, 22.6, 23.02.**

6. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt** having the following X-ray powder diffraction data : (2 θ): 18.28, 18.8, 19.8, 20.12, 20.62, 21.10, 21.44, 21.88, 22.6, 23.02; having a DSC exotherm at 194.60°C (onset at 185.28°C) and an endotherm at 245.81°C (onset at 226.23°C), and a water content of between 0.4 and 0.7% by weight as determined by titration according to Karl Fischer.
7. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt** according to claim 3, having an DSC endotherm at 242.18°C (onset at 207.85°).
8. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt** according to claim 5, having a DSC exotherm at 194.60°C (onset at 185.28°C) and an endotherm at 245.81°C (onset at 226.23°C).
9. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt** according to claim 3, wherein the solubility in a solution of pH 9.5 is 4.0 to 6.0 mg/ml.
10. The **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid arginine salt** according to claim 4, wherein the solubility in a solution of pH 9.5 is 4.0 to 6.0 mg/ml.
11. **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt** according to claim 5, wherein the solubility in a solution of pH 9.5 is 38.0 to 40.0 mg/ml.
12. **S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt** according to claim 6, wherein the solubility in a solution of pH 9.5 is 38.0 to 40.0 mg/ml.